

THE NEARBY OPTICAL GALAXY SAMPLE: THE LOCAL GALAXY LUMINOSITY FUNCTION

Christian Marinoni¹, Pierluigi Monaco^{1,3}, Giuliano Giuricin^{1,2}
and **Barbara Costantini^{1,2}**

¹ Dipartimento di Astronomia, Università di Trieste,
via Tiepolo 11, 34131 Trieste, Italia.

² SISSA, via Beirut 4, 34013 - Trieste, Italia.

³ Institute of Astronomy, Madingley Road, Cambridge, CB3 0HA, UK.

ABSTRACT

In this paper we derive the galaxy luminosity function from the Nearby Optical Galaxy (NOG) sample, which is a nearly complete, magnitude-limited ($B \leq 14$ mag), all-sky sample of nearby optical galaxies (~ 6400 galaxies with $cz < 5500$ km s $^{-1}$). For this *local* sample, we use galaxy distance estimates based on different peculiar velocity models. Therefore, the derivation of the luminosity function is carried out using the locations of field and grouped galaxies in real distance space.

The local field galaxy luminosity function in the B system is well described by a Schechter function with a slope $\alpha \sim -1.1$, a low normalization factor ($\Phi^* \sim 0.006$ Mpc $^{-3}$), and a particularly bright characteristic magnitude ($M_B^* \sim -20.6$) ($H_0 = 75$ km $^{-1}$ Mpc $^{-1}$). The exact values of the Schechter parameters slightly depend on the adopted peculiar velocity field models. Peculiar motion effects are of the order of statistical errors and cause at most variations of 0.8 in α and 0.2 mag in M_B^* . Our M_B^* -value is brighter by a few tenths of magnitude than previous corresponding values, because, referring to total corrected blue magnitudes, better represent the galaxy light.

Also the selection function, evaluated in terms of the luminosity function, appears to be little sensitive to the adopted peculiar velocity field models, which, however, bear a large impact on the local galaxy density on the smallest scales.

The shape of the luminosity function of spiral galaxies does not differ significantly from that of E-S0 galaxies. On the other hand, the late-type spirals and irregulars have a very steeply rising luminosity function towards the faint end ($\alpha \sim -2.3 - -2.4$), whereas the ellipticals appreciably decrease in number towards low luminosities.

The presence of galaxy systems in the NOG sample does not affect significantly the field galaxy luminosity function, since environmental effects on the total luminosity function appear to be marginal. The luminosity function of the members of the richest galaxy systems tends to show a slightly brighter M_B^* -value than the norm.

In the light of constraints imposed by the observed galaxy number counts, the low normalization of the luminosity function suggests that the nearby universe ($cz \lesssim 5000$ km/s) examined in this paper may be underdense by a factor ~ 1.5 .

Subject headings: galaxies: distances and redshifts – galaxies: fundamental parameters – galaxies: luminosity function, mass function – cosmology: observations

1 INTRODUCTION

Much work has gone into the determination of the local ($z = 0$) optical galaxy luminosity function (LF) (see, e.g., the review by Binggeli, Sandage & Tammann 1988 for early works), which is a fundamental quantity in cosmology. However, there is still uncertainty in its normalization and in its faint end. These quantities are important for the question of whether there is significant evolution in galaxy properties to $z \sim 0.5$ and for the explanation of the faint galaxy counts. As stressed in the reviews by Colless (1997) and Ellis (1997), many efforts are still needed to verify the detailed form of the local LF. Furthermore, in a magnitude-limited galaxy sample the knowledge of the LF is an essential ingredient for evaluating the selection function for the sample.

In this paper we determine the optical galaxy LF —and, hence the intimately related selection function — of a magnitude-limited complete sample of nearby optical galaxies. The present work is indeed meant to be an important step towards the recovering of the optical galaxy density field in the nearby universe (see, e.g., Marinoni et al. 1998a for a preliminary account of this work). As previously discussed (e.g., Hudson 1993, Santiago et al. 1995), optical galaxy samples are more suitable for mapping the galaxy density field on small scales than IRAS-selected samples, which have been frequently used as tracers of the galaxy density field on large scales.

In this work we consider the magnitude-limited, all-sky sample of nearby galaxies which was extracted by Garcia et al. (1993) from the Lyon-Meudon Extragalactic Database (LEDa). This sample comprises 6392 galaxies with recession velocities $cz < 5500 \text{ km s}^{-1}$ and corrected total blue magnitudes $B \leq 14 \text{ mag}$. Although the different optical galaxy catalogues, from which data are collected and homogenized in the LEDa, have different limits of completeness in apparent magnitude or angular diameter, the above-mentioned galaxy sample was found to be substantially complete down to its limiting magnitude $B = 14 \text{ mag}$, for galactic latitudes $|b| > 20^\circ$ (Garcia et al. 1993). On the basis of the LEDa compilation of bright galaxies with and without redshifts, we estimate that the completeness level of this sample limited to $|b| > 20^\circ$ (5366 galaxies) is $\sim 85\%$. For the sample of 5832 galaxies having $|b| > 15^\circ$, the completeness level decreases to $\sim 80\%$, according to the deficit of galaxy counts at low $|b|$ -values. Most the bright galaxies with unknown redshift lie at $|b| < 30^\circ$.

Group assignments for the galaxies of this sample have been already provided by Garcia (1993), who identified groups by employing the percolation or *friends of friends* method proposed by Huchra & Geller (1982) and the hierarchical clustering method (e.g., Tully 1987). The adopted final catalog of groups was defined as that one which includes only groups common to the two catalogs. Accordingly, we shall treat 3381 ungrouped galaxies as field galaxies. Of the 3011

members of the 485 groups (with at least three members), in the present study we omit 22 galaxies which are possible members (members which almost satisfy the selection criteria).

Garcia et al. (1993) tabulated several parameters for each galaxy, such as the morphological type, the corrected total blue magnitude B and the corrected angular size (the latter transformed in the standard systems of the RC3 catalog by de Vaucouleurs et al. 1991), and the distance modulus. In general, this quantity has been derived simply from redshift-dependent distances with an adopted Hubble constant $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$, except for the nearest galaxies (with $cz < 1500 \text{ km s}^{-1}$), where redshift-independent distances obtained from some distance indicators (DIs) (i.e. from a blue Tully-Fisher relation which relates luminosities to maximum rotation velocities and a luminosity-luminosity index relation) have been also partially considered.

In Paper I (Marinoni et al. 1998b), we corrected the redshift-distances of the galaxies and groups of this sample for peculiar motions by means of models of the peculiar velocity field. We purposely avoided correcting distances by calculating the peculiar velocity field on the basis of the positions and redshifts of the galaxy sample itself (in the linear theory approximation), because the galaxy sample does not presumably include all relevant gravitational sources (such as the Shapley concentration) for local peculiar motions. We instead followed two basic independent approaches to modeling the velocity field in the nearby universe: i) a modified cluster dipole model, which is Branchini & Plionis' (1996) optical cluster dipole reconstruction scheme modified by the inclusion of a local model of the Virgocentric infall in the Local Supercluster region. ii) a semi-linear approach which uses a multi-attractor model fitted to the Mark II and Mark III catalogs of galaxy peculiar velocity (Willick et al. 1997). In this model the velocity field was assumed to be generated by a few prominent gravitational sources (Virgo cluster, Great Attractor, Perseus-Pisces and Shapley superclusters), which were described as spherically symmetric masses with King-type mass density radial distributions.

Furthermore, we inverted the redshift-distance relations relative to different velocity field models and solved the problem of the triple-valued zones of this relation (these zones can appear in the vicinity of prominent overdensities) by using blue Tully-Fisher relations calibrated on suitably defined samples of objects having distances predicted by peculiar velocity models.

In this way, in Paper I we provided homogeneous estimates of distances for the individual 3381 field galaxies and the 485 groups of the above-mentioned galaxy sample, hereinafter denoted as Nearby Optical Galaxy (NOG) sample.

There are three main conceptual differences between our present analysis of the local galaxy LF and previous relevant studies based on other, comparably shallow samples, such as the CfA1

(Davis & Huchra 1982), CfA2 (Marzke et al. 1994a), SSRS2 (da Costa et al. 1994), and ORS (Santiago et al. 1995, 1996).

First, we use the distance estimates given in Paper I for determining the galaxy LF of the NOG sample. Thus, our derivation of the galaxy LF is carried out using the locations of field and grouped objects in real distance space, not in redshift space as is usually done in the literature. Having calculated different models of galaxy distances, we are also able to quantify to what extent differences in the current views on the cosmic flows can affect the determination of the galaxy LF of an optical galaxy sample. We will find that the luminosity and selection function of our optical and local sample are little sensitive to the adopted peculiar velocity field, thus confirming Yahil et al.'s (1991) finding based on IRAS galaxy samples (and different velocity field models).

Second, we rely on apparent total B magnitudes fully corrected for internal extinction, Galactic extinction and K-dimming (Garcia et al. 1993). These corrections lead to bright magnitudes. In particular, the first correction, which is conspicuous in very inclined spiral galaxies, is generally neglected in generic redshift surveys, which comprise many faint galaxies of unknown inclination. The average correction for generic samples of bright spirals is ~ 0.2 – 0.3 mag in the B band (de Vaucouleurs, de Vaucouleurs & Corwin 1976; Hasegawa & Umemura 1993). Being selected on the basis of homogenized corrected magnitudes, NOG is in principle designed to provide a good estimate of the bright end of the local LF.

Third, the NOG sample, which contains a large catalog of groups selected in a homogeneous way according to well-defined selection criteria, allows us to investigate on possible environmental effects on the LF, specifically on differences between the LFs of non-grouped (field) and grouped galaxies.

A different, complementary approach to the definition of an all-sky sample of nearby optical galaxies with good completeness in redshift, the "Optical Redshift Survey" (ORS), was followed by Santiago et al. (1995). The ORS sample contains 8286 galaxies with known redshift and consists of two overlapping optically-selected samples (limited in apparent magnitude and diameter, respectively) which cover almost all the sky with $|b| > 20^\circ$. Each sample is a concatenation of three subsamples drawn from the Uppsala General Catalogue (UGC) in the north, the European Southern Observatory (ESO) catalogue in the south, and the Extension to the Southern Observatory Catalogue (ESGC) in a strip just south of the celestial equator. The authors selected their own galaxy sample according to the raw (observed) magnitudes and diameters and then quantified the effects of Galactic extinction (as well as random and systematic errors) on the resulting galaxy density field, which was calculated out to $cz=8000$ km/s in redshift space

(Santiago et al. 1996). Adding the IRAS 1.2 Jy galaxy sample (Fisher et al. 1995) in the unsurveyed zone of avoidance ($|b| > 20^\circ$) and at large distances ($cz > 8000$ km/s), Baker et al. (1998) calculated the resulting peculiar velocity field.

Notwithstanding the differences in selection criteria and the larger redshift incompleteness of the NOG, the distribution of NOG galaxies on the sky is qualitatively similar to that of ORS galaxies; both samples delineate similar major structures in the nearby universe.

The outline of our paper is as follows. In §2 we address the determination of the Schechter-type blue LF for the NOG sample. In §3 we describe our results concerning the total LF, the morphological type-specific LFs, and the comparison between the field and grouped galaxy LFs. In §4 we compare our results with previous LF determinations and discuss some implications related to galaxy number counts. In §5 we define the selection function of the NOG sample in terms of the galaxy LF. Conclusions are drawn in §6.

Throughout, the Hubble constant is $75 \ h_{75} \ km \ s^{-1} Mpc^{-1}$.

2 The galaxy luminosity function: analysis

We evaluate the galaxy LF of the NOG sample in the standard B system of the RC3 catalog. We rely on the various sets of distances (corrected for peculiar motions) given by Marinoni et al. (1998b) for field and grouped galaxies and on the total B magnitudes as derived by Garcia et al. (1993), who transformed the original raw data to the standard B system of the RC3 catalog. For numerous NOG galaxies total B magnitudes were carefully derived from aperture photometry and detailed (photographic or CCD) surface photometry. Garcia et al. (1993) corrected the total B magnitudes for internal extinction (following Tully & Fouqué 1985), Galactic extinction (following Burstein & Heiles 1978 a,b, 1982, 1984), and K-dimming (Pence 1976).

In general, extensive, bright magnitude-limited samples such as the NOG sample, which is currently one of the largest samples used in the LF determination, can provide good determinations of the local LF, except for its faint end (i.e., roughly for $M_B > -15$). Compared to other wide-angle, comparably shallow, bright magnitude-limited galaxy samples, such as the CfA1 (Davis & Huchra 1982), CfA2 (Marzke et al. 1994a), and SSRS2 (da Costa et al. 1994; Marzke & da Costa 1997) redshift surveys, the NOG sample may be less sensitive to local density fluctuations because it covers a much larger solid angle.

Deeper magnitude-limited redshift surveys are in principle better designed to address issues concerning the faint end of the local LF and its normalization, which in shallow galaxy samples could be affected by local density fluctuations. But, in practice, surveys which extend local samples to fainter magnitudes and higher redshifts may begin suffering more seriously from

some kinds of uncertainties: i) their photometric quality gets worse and, in particular, the photographic photometry scale may be not well-defined (see, e.g., Takamiya, Kron & Kron 1995, Metcalfe, Fong & Shanks 1995, Rousseau et al. 1996, for systematic errors in Zwicky magnitudes, APM and COSMOS bj photometry, respectively); ii) their completeness gets more doubtful, because, at a given magnitude, they tend to miss more compact and low-surface brightness (LSB) galaxies by virtue of selection effects inherent in standard image detection algorithms (e.g., Disney & Philipps 1985; Davies, Disney & Phillipps 1989). As a matter of fact, at a given magnitude, galaxies which lie at nearer distances are selected over a broader interval of surface brightness (e.g., Phillipps, Davies, & Disney 1990).

Much of the meaningful discrepancies between the local LFs derived in the literature (see §4 below) can be ascribed to photometric problems (magnitude scale errors, use of different isophotal or aperture magnitude systems, lack of adequate corrections of observed magnitudes) and to different observational selection criteria used in constructing galaxy samples. Both kinds of problems are a little minimized in NOG, because this sample comprises both bright and nearby galaxies.

Furthermore, the deepest surveys suffer from redshift dependence in the measured isophotal or aperture magnitudes, due to the combined effects of the point spread function and surface brightness variation (e.g., Dalcanton 1998). In any case, very faint magnitude-limited redshift surveys (e.g., Ellis 1997), which are very important tools for probing evolutionary effects on the galaxy LF, have difficulties in giving unambiguous constraints on the local galaxy LF, because they are sensitive to evolutionary effects (e.g., the problem of the faint blue galaxy excess) and to the local normalization problem (e.g., Driver & Phillipps 1996).

2.1 Estimating the shape

We evaluate the shape of the galaxy LF following Turner's (1979) method (see also de Laparent, Geller & Hucra 1989). As recently demonstrated by Willmer (1997), who compared the robustness of different Schechter-type LF estimators by means of Monte Carlo simulations, this method — as well as all widely used methods, except the $1/V_{max}$ technique — gives a substantially unbiased estimate of the shape of this function, especially in the case of large samples.

Under the assumption that the LF is universal (i.e. independent of location in the universe), the galaxy number per unit volume and magnitude has a form separable into a product of functions of absolute magnitude M and position $\Phi(\mathbf{r}, M)dMdr^3 = \varphi(M)\rho(\mathbf{r})dMdr^3$.

In this notation, φ represents the fraction of objects with luminosity M , i.e. the shape of LF.

For a magnitude-limited sample of galaxies, the mean number of objects with luminosity M is

$$n(M)dM = \left[\int_{V_{max}(M)} \rho(\mathbf{r})dV \right] \varphi(M)dM = S(M)\varphi(M)dM \quad (1)$$

where $V_{max}(M)$ is the maximum volume in which galaxies brighter than M are still visible.

Considering the number density of galaxies brighter than M

$$N(< M) = S(M) \int_{-\infty}^M \varphi(M')dM' = S(M)\Psi(M) \quad (2)$$

and taking the ratio between eq. 1 and eq. 2, we are left with a quantity $Y(M)dM$ which is density-independent. Rewriting this quantity using the following differential form

$$Y(M)dM = \frac{d\Psi}{\Psi(M)}, \quad (3)$$

an analytic expression for the shape of the LF is given by

$$\varphi(M) = \phi_0 Y(M) \exp \int_{M_0}^M Y(M')dM' \quad (4)$$

where M_0 is the brightest absolute magnitude we use in the estimation of the shape of the LF.

Inside $V_{max}(M)$, and for small absolute magnitude increments (i.e. for small distance increments δr), $Y(M)$ can be estimated using the following approximation (de Lapparent, Geller & Huchra, 1989)

$$Y(M) \approx \frac{N(\leq M) - N(\leq M - \delta M)}{N(\leq M)\delta M} \quad (5)$$

where $N(\leq M)$ is the number of galaxies brighter than M . Even if this estimation technique is intrinsically non parametric, in order to make comparisons with previous works, and also in order to avoid biases in the determination of the LF shape (de Lapparent et al., 1989), we fit eq. 5 against the one predicted, assuming, for the LF shape φ in the $N(<M)$ estimator (eq. 2), the Schechter (1976) form

$$\varphi \propto 10^{0.4(M^* - M)(\alpha + 1)} \exp[-10^{0.4(M^* - M)}]. \quad (6)$$

In this equation, M^* is a fiducial magnitude which characterizes the point (the "knee" of the function) where the form of the function changes from a power-law (with slope α) to an exponential slope

With this least-square algorithm, weighted assuming Poisson fluctuations, we can measure the best parameters α and M^* for the Schechter function.

However, in order to account for random errors in the magnitudes, we convolve the Schechter function $\Phi_s(M)$ with a Gaussian magnitude error distribution with zero mean and dispersion $\sigma_m = 0.2$ mag to give an observed LF $\Phi_o(M)$:

$$\Phi_o(M) = \frac{1}{\sqrt{2\pi}\sigma_m} \int_{-\infty}^{+\infty} \Phi_s(M') \exp\left[-\frac{(M' - M)^2}{2\sigma_m^2}\right] dM'. \quad (7)$$

We perform the least-square fit of the LF estimator Y using various distance intervals of $\delta r = 250, 500$ and 700 km/s. The lower limit for δr was fixed by the spread caused by random noise in the peculiar velocity field, while the upper limit was set at the peak value of the absolute strength of the peculiar velocities. A variation in δr over the range $250\text{--}700$ km s^{−1} causes changes in the LF parameters α and M^* which are of the order of uncertainties.

2.2 Estimating the normalization

Involving ratios between the differential and the integrated LFs, the estimator Y does not provide any information about the normalization factor ϕ^* (i.e., the scaled number density of galaxies with luminosity M^*) which has to be derived independently. We normalize the Schechter function using the fact that the mean spatial galaxy density n is related to the LF by the relation

$$\phi^* = \frac{n}{\int_{-\infty}^{M_s} \Phi(M) dM}, \quad (8)$$

where $\Phi_o(M)$ is the convolved Schechter function with ϕ^* set equal to one. Since the LF is poorly constrained at the faint end, we cut off the integral at the limit of $L_s = L_{min}(r_s)$, where $L_{min}(r)$ is the minimum luminosity necessary for a galaxy at distance r (in Mpc) to make it into the sample and r_s is taken equal to $r_s = 500/(75 \cdot h_{75}) \sim 6.7 h_{75}^{-1}$ Mpc. $L_{min}(r)$ corresponds to the absolute magnitude $M_B = -5 \log r - 25 + B_{lim}$, where $B_{lim} = 14$ mag is the limiting apparent magnitude of our galaxy sample; thus L_s corresponds to $M_s = -15.12 + 5 \log h_{75}$.

Various methods have been proposed to estimate the mean density. Given our complete and magnitude-limited sample of galaxy with $r_{min} < r_i < r_{max}$, we calculate the mean galaxy density n using the unbiased estimator

$$fn = \frac{\sum_i^{N_{gal}} w(r_i)}{\int_{r_{min}}^{r_{max}} \left(\frac{dV}{dr}\right) dr S(r) w(r)}, \quad (9)$$

where $S(r)$ is the selection function given in equation 11 below, $f=0.8$ is the adopted com-

pleteness level (see §1) and $w(r)$ is the weight function

$$w(r) = \frac{1}{[1 + 4\pi n J_3(r_c) F(r)]}, \quad J_3(r_c) = \int_0^{r_c} r^2 \xi(r) dr \quad (10)$$

which minimizes the variance in the estimate of n (Davis & Huchra, 1982). For the second moment of the spatial two-point galaxy correlation function $\xi(r)$, we have adopted the value $J_3 = 320h^{-3} \text{Mpc}^3$ obtained for $r_c = 40 h^{-1} \text{Mpc}$ (Marzke et al. 1994a). Even if the value of J_3 is uncertain, because the contribution for $r > r_c$ could be substantial, the final result for n (and ϕ^*) depends little on the value of J_3 used (Loveday et al. 1992). In our case, doubling or halving the adopted value of J_3 makes a few per cent difference to n .

In this minimum-variance weighting scheme, n is determined iteratively using eqs. 9 and 10 while, for errors estimate, we add in quadrature the uncertainties in n arising from galaxy clustering ($\delta n \sim n[4\pi J_3/V]^{0.5}$) with those arising from varying the parameters of the luminosity function along their joint 1σ error ellipse (Lin et al. 1996).

We have checked that the incompleteness of NOG does not appreciably depend on galaxy morphological type and mean surface brightness (inside the isophote 25 B-mag arcsec⁻²), whereas it gets weaker at brighter B magnitudes (e.g., $f=0.90$ (0.92) for $|b| > 15^\circ$ ($|b| > 20^\circ$) and $B \leq 13.5$; $f=0.93$ (0.96) for $|b| > 15^\circ$ ($|b| > 20^\circ$) and $B \leq 13.0$). After having derived in detail the degree of completeness of NOG as a function of the B magnitude, under the simple assumption of a uniform distribution of the missing galaxies in redshift space, we have calculated that this dependence does not appreciably affect the LF shape; it translates into a bias towards M^* -values systematically brighter by 0.1 mag (and to α -values systematically flatter by 0.01 at most).

In order to take into account border effects which arise from the mapping between redshift space and true-distance space and which influence the completeness of the sample (see Fig. 2 below), we evaluate the Schechter-type LF for the galaxies with distances $r < 5250/(75 h_{75}) \text{Mpc}$ only. Further we consider only galaxies having $|b| > 15^\circ$, in order to avoid a large incompleteness, and M_B -values in the range $-22.5 \leq M_B - 5 \log h_{75} \leq M_s$ (where $M_s = -15.12$), in order to avoid small number statistics in each bin.

We use the sets of distances r relative to different peculiar velocity field models, i.e., the multi-attractor model based on the whole Mark III data and on Mark III spiral data (Mark III*), Branchini & Plionis' (1996) cluster dipole model, the cluster dipole model as modified by Marinoni et al. (1998b) with the inclusion of a local model of the Virgocentric infall, the unperturbed Hubble flows in the CMB and Local Group (LG) frames. The number of galaxies ($N \sim 5350$) which fulfills the above-mentioned selection criteria changes according to the distance model adopted.

3 The galaxy luminosity function: results

3.1 The total luminosity function

Table 1 reports the Schechter parameters obtained for sets of distance models. The errors given for α and M_B^* are the projections onto their axes of the 1σ confidence ellipse level, as derived from the χ^2 matrices of the least-square fit of $Y(M_B)$.

The 1σ formal errors in ϕ^* include contributions both from galaxy clustering and from uncertainties in M_B^* and α . We have checked that neglecting magnitude errors makes the slope α steeper by 0.03 and M^* brighter by 0.1 mag, on average.

The last columns of Table 1 contain the galaxy luminosity density $\rho_L = \int_{L_s}^{\infty} L\Phi(L)dL$ and the galaxy number density n together with the 1σ errors. Letting the lower integration limit go to zero does not increase ρ_L significantly.

Fig. 1 show our results for the normalized LFs and the 1σ error ellipses in the (α, M_B) plane, for galaxy distances predicted by the Mark III multi-attractor model and the cluster dipole model. Fig. 1 and the values of χ^2/dof given in Table 1 indicate that the Schechter function always gives a good fit to the LF of our sample. In the lowest luminosity bin there might be a hint for an excess of objects relative to the plotted fits for the Mark III multi-attractor models, but not for the cluster dipole model.

Fig. 2 displays the observed galaxy number-distance histograms together with the curves which show the expected distribution for an uniform universe, for different distance models.

TABLE 1

THE PARAMETERS OF THE SCHECHTER LFs FOR DIFFERENT PECULIAR VELOCITY FIELD MODELS.

Model	α	$M_B^* - 5 \log h_{75}$	ϕ^*	χ^2/dof	ρ_L	n
			$(10^{-3} h_{75}^3 Mpc^{-3})$		$(10^8 L_{\odot} h_{75} Mpc^{-3})$	$(10^{-2} h_{75}^3 Mpc^{-3})$
Multi-attractor (Mark III)	-1.10 ± 0.06	-20.67 ± 0.08	5.9 ± 0.9	0.52	1.78 ± 0.28	3.2 ± 0.6
Multi-attractor (Mark III*)	-1.07 ± 0.06	-20.70 ± 0.08	5.6 ± 0.9	0.42	1.67 ± 0.29	2.9 ± 0.5
Cluster dipole model	-1.15 ± 0.05	-20.63 ± 0.08	5.9 ± 0.9	0.54	1.75 ± 0.30	3.7 ± 0.8
Modified cluster dipole model	-1.09 ± 0.07	-20.82 ± 0.08	5.8 ± 0.9	0.68	1.76 ± 0.33	3.2 ± 0.7
Hubble Flow (CMB)	-1.11 ± 0.06	-20.63 ± 0.07	6.2 ± 1.0	0.49	1.80 ± 0.29	3.5 ± 0.7
Hubble Flow (LG)	-1.15 ± 0.05	-20.64 ± 0.08	5.9 ± 0.9	0.24	1.78 ± 0.31	3.7 ± 0.8




Fig. 1.— We plot the normalized differential LFs (together with 1σ error bars) for galaxy distances derived from the multi-attractor model based on all Mark III data (*left*) and from the cluster dipole model data (*right*).




Fig. 2.— The plots show the observed galaxy number-distance histograms. They refer to redshift-distances obtained using the Hubble relation in the CMB frame (a) and in the LG frame (b), and to distances corrected according to the multi-attractor model fitted to the Mark III data (c), the multi-attractor model fitted to the Mark III* data (d), the cluster dipole model (e), and the modified cluster dipole model (f). The superimposed solid curves indicate the predictions given by the best Schechter function fits in the case of a homogeneous galaxy distribution.

The details of the large "peak and trough" fluctuations, which are seen both in wide-angle (e.g., the CfA2 sample) and narrow-angle surveys (e.g., the ESP sample by Zucca et al. 1997) are quite different in the various plots. Thus, the effects of peculiar motions have a large impact on the evaluation of the local galaxy density, especially on the smallest scales. On large scales, an overdensity at a distance of $\sim 1500\text{--}2000 \text{ km s}^{-1}$ (corresponding to the center of the Local Supercluster) and an underdensity around $\sim 3500 \text{ km s}^{-1}$ are common to all histograms.

All cases lead to similar values of ϕ^* , ρ_L , n , and α . All cases give also similar M_B^* -values, except for the modified cluster dipole model. Owing to its large Virgocentric infall, the modified cluster dipole model moves nearby galaxies quite close to us (see Fig. 2f). Moreover, it gives a M_B^* -value typically brighter by $\sim 0.1 - 0.2 \text{ mag}$ than the norm, in agreement with the discussion by Efstathiou, Ellis & Peterson (1988) on the effect of a comparable Virgocentric infall on the LF of the RSA galaxy sample (Sandage & Tammann 1981). However, in recent years there is a growing evidence that the Virgo cluster is not a major source of the peculiar velocity field in the Local Supercluster. From the analysis of the magnitudes of bright cluster galaxies in a way which is free of assumptions about motions in the LS, Gudehus (1995) found no significant evidence of a Virgo infall of the LG. Therefore, in what follows we do not longer use the modified cluster dipole model, which is likely to be the least realistic model.

To sum up, our calculations prove that differences in the peculiar velocity field models, though appreciably affecting the galaxy density field on galaxy-galaxy distance scale, do not have large effects on the optical LF of a shallow galaxy sample which covers a very large solid angle. The effects could be greater in samples restricted to fairly narrow solid angles (see, e.g., Davis & Huchra 1982, for the effect of a Virgocentric infall on the northern part of the CfA1 sample).

The values of ρ_L and n tabulated in Table 1 refer to the population of cataloged galaxies of highish surface brightness. In general, magnitude-limited redshift surveys miss an increasing fraction of LSB objects at fainter magnitudes, which makes it difficult to determine the faint end of the LF. Among specific investigations designed to recover the LSB field population (e.g., the review by Impey & Bothun 1997), the large APM LSB redshift survey allowed the reconstruction of the LF of LSBs (Sprayberry et al. 1997), which shows that the LSB population (dominated by late-type spirals and irregulars), becomes appreciably numerous for magnitudes fainter than $M_B \sim -15$ only.

Hence, missing LSB objects do not affect appreciably the bright end of our LF. In the faint end they contribute strongly to the number density of galaxies (presumably also to cosmological mass density in baryons; e.g., Bristow & Phillipps 1994), but not too much to the luminosity density. Sprayberry et al. (1997) evaluated a contribution of $0.2 h_{75} L_{\odot} \text{ Mpc}^{-3}$ (in the B

system) from the LF of the APM LSB galaxies, whereas Dalcanton et al. (1997a) derived $\rho_L = 0.1 h_{75} L_{\odot} \text{ Mpc}^{-3}$ (in the B system) from their small sample of late-type LSB galaxies identified in a large-area transit scan CCD data.

With the addition of the $\sim 10\%$ light contribution given by Sprayberry et al. (1997), our mean value of $\rho_L = 2.0 \pm 0.3 L_{\odot} h_{75} \text{ Mpc}^{-3}$ implies a critical mass-to-light ratio $(\frac{M}{L})_c = \frac{1.563 \cdot 10^{11} h_{75}}{\rho_L} = 7.8 \pm 1.2 \cdot 10^2 h_{75} (M/L)_{\odot}$ to close the universe. The cosmological density parameter $\Omega_0 = (M/L)/(M/L)_c$ is $\sim 0.2\text{--}0.3$ if we set $M/L \sim 200 h_{75} \cdot (M/L)_{\odot}$, a typical virial mass-to-light ratio estimated from galaxy clusters in the blue (e.g., Bahcall, Lubin & Dorman 1995; David, Jones & Forman 1995; Girardi et al. 1999). However, on larger scales, our analysis of cosmic flows (see Paper I) leads to a larger value of Ω_0 , $\Omega_0 \sim 0.5$ (see also the reviews by Dekel, Burstein & White 1997 and Hamilton 1998).

3.2 The type-specific luminosity functions

Lumping together giants and dwarfs, we divide the NOG galaxies into five type bins, the ellipticals (E), lenticulars (S0), the Sa–Sb spirals (which cover the range between S0/a and Sb), the Sc–Sd spirals (which cover the range between Sbc and Sd), and the Sm–Im objects. We also consider the division into two type bins, the E–S0 objects and the spirals and irregulars (S–Im). The morphological types are available for almost all the NOG galaxies.

In Table 2 we give the results for the type-specific LFs (over the same ranges of distances and absolute magnitudes used for the whole sample), for different sets of distance models. In Fig. 3 we plot the type-specific LFs and the total LF, in the specific case of Mark III distances. In Fig. 4 we plot the 1σ confidence ellipses for the parameters α and M_B^* relative to the LFs shown in Fig. 3 for different morphological types.

The morphological type-dependence of the LF is little sensitive to the distance models adopted. Assembling together the E and S0 galaxies, we obtain a E–S0 LF which remains almost flat, although the LF of the giant objects alone can be described by a Gaussian function (e.g., Binggeli et al. 1998; Biviano et al. 1995). More precisely, the LF of E galaxies appreciably declines towards faint luminosities (with $\alpha \sim -0.5$), whilst the LF of S0 objects does not.

The number of Sa–Sb types declines towards fainter magnitudes according to a slope ($\alpha \sim -0.6 - -0.8$) which is considerably flatter than the norm for spirals, whilst the Sm–Im objects, which are on average, fainter than the norm, always exhibit a very steep slope ($\alpha \sim -2.3 - -2.4$). The slope α tends to get progressively steeper as one goes from early spirals to late spirals and irregulars. The value of M_B^* is fainter for S0 galaxies than for the other types.

TABLE 2

THE PARAMETERS OF THE MORPHOLOGICAL TYPE-SPECIFIC LFs.

Model	Sample	Ngal	α	$M_B^* - 5 \log h_{75}$	$\phi^* (10^{-3} h_{75}^3 Mpc^{-3})$	χ^2/dof
Multi-attractor (Mark III)	E	344	-0.47 ± 0.22	-20.75 ± 0.26	0.46 ± 0.12	0.56
	S0	596	-1.17 ± 0.20	-20.44 ± 0.26	0.81 ± 0.20	0.37
	E-S0	940	-0.97 ± 0.14	-20.69 ± 0.18	1.03 ± 0.24	0.51
	Sa-Sb	1521	-0.62 ± 0.11	-20.51 ± 0.12	2.20 ± 0.46	0.34
	Sc-Sd	2240	-0.89 ± 0.10	-20.39 ± 0.11	3.12 ± 0.59	0.44
	Sm-Im	619	-2.41 ± 0.28	-21.11 ± 0.72	0.07 ± 0.07	0.68
	S-Im	4380	-1.10 ± 0.07	-20.63 ± 0.09	4.58 ± 0.73	0.63
Cluster Dipole Model	E	345	-0.56 ± 0.22	-20.71 ± 0.26	0.45 ± 0.12	0.39
	S0	605	-1.03 ± 0.21	-20.20 ± 0.24	1.03 ± 0.26	0.31
	E-S0	950	-1.03 ± 0.14	-20.67 ± 0.19	1.06 ± 0.25	0.64
	Sa-Sb	1563	-0.73 ± 0.11	-20.48 ± 0.12	2.24 ± 0.47	0.49
	Sc-Sd	2289	-0.97 ± 0.09	-20.35 ± 0.12	3.17 ± 0.60	1.09
	Sm-Im	597	-2.45 ± 0.32	-21.12 ± 0.73	0.07 ± 0.07	0.64
	S-Im	4449	-1.17 ± 0.08	-20.60 ± 0.09	4.52 ± 0.72	0.89
Hubble Flow in LG frame	E	346	-0.55 ± 0.22	-20.73 ± 0.27	0.45 ± 0.11	0.77
	S0	589	-1.08 ± 0.21	-20.18 ± 0.24	0.79 ± 0.20	0.27
	E-S0	935	-1.12 ± 0.13	-20.72 ± 0.19	0.95 ± 0.23	0.74
	Sa-Sb	1539	-0.78 ± 0.11	-20.57 ± 0.13	2.10 ± 0.44	0.57
	Sc-Sd	2249	-0.93 ± 0.09	-20.33 ± 0.11	3.21 ± 0.61	0.37
	Sm-Im	620	-2.27 ± 0.32	-20.75 ± 0.69	0.15 ± 0.15	0.42
	S-Im	4408	-1.17 ± 0.07	-20.60 ± 0.09	4.62 ± 0.70	0.45

Except for the earliest (E) and latest (Sm–Im) types, there are no large differences between the shapes of the type-specific LFs. In particular, the shape of the E–S0 LF does not differ significantly from that of the S–Im objects.




Fig. 3.— We plot the normalized LFs for all galaxy types and for different morphological types, adopting galaxy distances as given by the multi-attractor model fitted to the Mark III data.




Fig. 4.— We plot contours representing the 1σ deviation from the best-fitting values of α and M_B^* relative to the LFs shown in Fig. 3 for different morphological types.

3.3 The luminosity function of group galaxies

With the purpose of investigating on the effect of the presence of galaxy systems on the LF determination, we recalculate the LFs after removing the members of the systems containing more than 5, 10, and 20 galaxies (e.g., removing 114, 42, and 10 systems for a total of 1437, 830, and 355 members, respectively, in the specific case of the Mark III multi-attractor model). We never find any significant difference in α and M_B^* with respect to the values tabulated in Table 1 for the whole sample.

Furthermore, we calculate the LFs for galaxy members of groups and for field (non-grouped) galaxies, separately, according to the various distance models. Table 3 lists the results. We find no significant differences between the LF derived from the two subsamples, whenever, for constructing the sample of grouped galaxies, we take all systems or a subsample of systems dominated by groups with few members (e.g., all the systems with $n > 5$ members). For all sets of distance models, we detect some marginal systematic effects (at the $\sim 1\sigma$ confidence level) if we restrict ourselves to the richest systems containing $n \geq 10$, $n \geq 15$, $n \geq 20$ members.

TABLE 3

THE SCHECHTER LFs FOR FIELD AND GROUP GALAXIES.

Model	Sample	Ngal	α	$M^* - 5 \log h_{75}$	χ^2/dof
Multi-attractor (Mark III)	field	2652	-1.19 ± 0.10	-20.59 ± 0.12	0.23
	groups	2668	-1.02 ± 0.07	-20.77 ± 0.10	0.58
	groups($n \geq 10$)	963	-1.21 ± 0.11	-20.99 ± 0.18	0.58
	groups($n \geq 20$)	394	-1.28 ± 0.18	-21.00 ± 0.31	0.28
Cluster dipole model	field	2735	-1.24 ± 0.09	-20.57 ± 0.12	0.59
	groups	2664	-1.07 ± 0.08	-20.69 ± 0.11	0.38
	groups($n \geq 10$)	937	-1.20 ± 0.12	-20.95 ± 0.23	0.62
	groups($n \geq 20$)	416	-1.07 ± 0.17	-20.83 ± 0.28	0.49
Hubble Flow in LG frame	field	2637	-1.18 ± 0.09	-20.63 ± 0.12	0.25
	groups	2706	-1.15 ± 0.07	-20.70 ± 0.11	0.43
	groups($n \geq 10$)	952	-1.35 ± 0.11	-20.97 ± 0.19	1.06
	groups($n \geq 20$)	395	-1.37 ± 0.17	-21.05 ± 0.44	0.49

TABLE 4

THE PARAMETERS OF THE LFs FROM VARIOUS SAMPLES.

Sample	Reference	Ngal m_{lim}	$M_{min} - 5 \log h_{75}$	α	$M_B^* - 5 \log h_{75}$	$\phi^*(h_{75}^3 Mpc^{-3})$
Stromlo-APM	Loveday et al., 1992	~ 1700 $b_j = 17.15$	-15.9	-0.97 ± 0.15	-20.42 ± 0.13	$(0.59 \pm 0.07) \cdot 10^{-2}$
CfA2	Marzke et al., 1994	~ 9100 $m_z = 15.5$	-17.9	-1.0 ± 0.2	-20.2 ± 0.3	$(1.7 \pm 0.4) \cdot 10^{-2}$
LCRS	Lin et al., 1996	~ 18700 Gunn $r \sim 17.4$	-17.4	-0.70 ± 0.05	-20.11 ± 0.02	$(0.8 \pm 0.04) \cdot 10^{-2}$
Autofib	Ellis et al., 1996	~ 600 $b_j = 24$	-14.9	-1.16 ± 0.05	$-20.22^{+0.15}_{-0.12}$	$(1.04^{+0.16}_{-0.13}) \cdot 10^{-2}$
SSRS2	Marzke & da Costa, 1997	~ 3300 $B(0) = 15.5$	-14.9	$-1.16^{+0.08}_{-0.06}$	-20.37 ± 0.08	$(0.46 \pm 0.13) \cdot 10^{-2}$
ESP	Zucca et al., 1997	~ 3300 $b_j = 19.4$	-13.3	$-1.22^{+0.06}_{-0.07}$	$-20.53^{+0.06}_{-0.08}$	$(0.8 \pm 0.02) \cdot 10^{-2}$
CS	Geller et al., 1997	~ 1700 $R = 16.13$	-15.6	-1.17 ± 0.19	$-20.35^{+0.17}_{-0.18}$	$(1.1 \pm 0.3) \cdot 10^{-2}$
Durham/UKST	Ratcliffe et al., 1998	~ 2100 $b_j \sim 17$	-15.9	-1.04 ± 0.08	-20.60 ± 0.10	$(0.72 \pm 0.01) \cdot 10^{-2}$

The LF of their members tend to show somewhat brighter values of M_B^* with respect to the field value, with no systematic shift of α . Although the average number of members of a group tends to decrease with increasing distance, the tendency is not so strong to affect significantly the above-mentioned results.

4 Discussion

4.1 Comparison with previous total galaxy luminosity functions

In Table 4 we show the Schechter function parameters of some recent redshift surveys of field galaxies. All published values have been reduced to $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$. We quote the original photometric passband of the survey, but we transform the characteristic absolute magnitude M^* to M_B^* through simple magnitude offsets; namely $b_j - r = +1.1$, which is the mean rest-frame color of the Las Campanas redshift survey (LCRS) galaxies (Lin et al. 1996), $b_j - R = +1.3$, which is the median rest-frame color of the Century Survey (CS) galaxies (e.g., Buta & Williams 1995; Geller et al. 1997), $b_j - m_Z = -0.45$ (the shift of the blue Zwicky magnitudes m_Z comes from the comparison of galaxy number-magnitude counts in Shanks et al. 1984); $B - b_j = -0.3$ as given by Efstathiou et al. 1988; $B - B(0) = -0.3$ (the zero point of the scale of the $B(0)$ magnitudes agrees with the b_j one within 0.2 mag, according to da Costa et al. 1994). The formal errors given in Table 4 do not include uncertainties related to color transformations. In the following, we quote the results given in the literature as scaled into the color and H_0 we are using.




Fig. 5.— This figure shows a comparison between the NOG LF we calculate in the case of the multi-attractor model based on Mark III data, and previous LFs obtained in the literature from recent redshift surveys of low depth (a) and intermediate depth (b).

Fig. 5 illustrates the comparison of our normalized LF (relative to the case of Mark III-based distances) with those obtained from some low-depth (Fig. 5a) and intermediate-depth (Fig. 5b) redshift surveys of field galaxies.

An accurate comparison between the LFs is impeded by the fact that applying simple constant magnitude offsets does neglect more complicated effects, which are due for example to galaxy colors, morphologies, and surface brightnesses. However, among the LFs there are certainly meaningful differences which are due to problems mentioned in §2. In any case, in the optical and near-infrared surveys mentioned in this section there is no indication of a systematic flattening of

the slope α , a simultaneous brightening of M_B^* , and a progressive decrease of the normalization factor with brighter surface brightness limits of the survey, as expected in the case in which observational selection effects depending on brightness surface limits were the dominant effects (e.g., Dalcanton, Spergel & Summers 1997b).

The M_B^* -values given in Table 4 are always appreciably fainter (by ~ 0.1 – 0.6 mag) than our values, which give a slower decline of the number of luminous galaxies with increasing luminosity.

Our α -values well agree with the average results obtained from the shallow surveys, such as the Stromlo-APM, CfA2, SSRS2, CS, Durham/UKST surveys, and from intermediate-depth redshift surveys, such as the ESP and Autofib survey (restricted to $0.02 < z < 0.15$). The only very discrepant value of α is that coming from LCRS. Its declining faint-end slope ($\alpha \sim -0.7$) can not be due to the fact that galaxies are selected in the red band, because the R-selected CS sample yields a much steeper slope. It can be partly related to the fact that a Schechter form does not fit very well the LCRS data around $M_B \sim -19.5$ and in the faint end (see Bromley et al. 1998) and mostly to the fact that LCRS may be biased against observing low-luminosity galaxies, because of the survey selection criteria (Geller et al. 1997).

In the intermediate luminosity interval ($M_B \sim -20$), our results point to a fairly low amplitude for the LF (see Fig. 5), in substantial agreement with the results coming from several shallow redshift surveys, especially with the Durham/UKST survey, which has yielded the LF which most closely resembles our results.

It is interesting, albeit less easy, to compare our results with those coming from the ORS. As a matter of fact, as warned by Santiago et al. (1996), it is difficult to properly compare the ORS LFs with other published LFs in the literature (including our results), because the ORS LFs are not corrected for Malmquist bias. Fig. 6 shows the comparison of our normalized LF (relative to the case of Hubble-flow distances in the LG frame) with the ORS LFs directly obtained from the UGC and ESO subsamples by Santiago et al. (1996), who parameterized the ORS LFs through a generalization of the Schechter form. For this comparison we apply no magnitude offset (since the ESO B system does not deviate too much from the standard RC3 B system; see e.g., Paturel, Bottinelli & Gouguenheim 1994), and no correction for differences in the mean galaxy density between the UGC and ESO subsamples. The NOG LF appears to be in satisfactory agreement with the ORS LFs at intermediate and faint magnitudes, whereas the bright end of the former seems to be shifted brightwards (like in the case of Fig. 6), suggesting magnitude offsets of ~ 0.5 and ~ 0.7 mag, for the ESO and UGC subsamples respectively. Notably, the latter offset (which refers to Zwicky magnitudes) is much smaller than the amount (~ 1.2 – 1.3 mag) required to match the NOG LF and the bright end of the CfA2 LF (which also uses Zwicky photometry).




Fig. 6.— This figure shows a comparison between the NOG LF that we calculate in the case of redshift-distances obtained using the Hubble relation in the Local Group frame and the ORS LFs obtained directly from the ESO and UGC subsamples of the ORS.

The anomalously high value of the CfA2 normalization is likely to be due to strong local inhomogeneities (e.g., the "Great Wall"). The quoted CS normalization, being the average value for the velocity range between 18 000 and 28 000 km s^{-1} , is also high, because in this range this survey slices through a portion of the Corona Borealis supercluster. Remarkably, both CfA2 and CS evidence that ϕ^* increases by a factor ~ 2 from nearby regions ($cz < 5000 \text{ km s}^{-1}$) to distant regions ($cz \sim 10000 \text{ km s}^{-1}$ and $\sim 25000 \text{ km s}^{-1}$), thus admitting a normalization consistent with our value in the nearby regions.

The deeper redshift surveys (ESP, LCRS, Autofib) disagree in the amplitude of the LF at

intermediate and faint magnitudes. Our results are intermediate with respect to the low LCRS value and the high values of the ESP and Autofib samples. Zucca et al. (1997) stressed the possibility of reconciling the high ESP normalization with the low values given by other surveys, by noting that the nearby region ($cz < 14000 \text{ km s}^{-1}$) encompassed by the ESP is underdense and that a recalculation of the LF for a subsample of bright ESP objects would give results consistent with those derived from the Stromlo-APM survey. This view seems to be qualitatively supported by the preliminary results of the intermediate-depth 2dF redshift survey (whose galaxies are selected from the same catalog used in the last survey); also this LF hints at a greater normalization than the Stromlo-APM one (Maddox 1998).

Noticeably, also the small-field deep near-infrared by Glazebrook et al. (1995b) and Cowie et al. (1996) disagree in the K-band LF normalization. The normalization found from the latter, larger survey is close to that coming from shallow near-infrared surveys (Mobasher, Sharples & Ellis 1993; Gardner et al. 1997; Szokoly et al. 1998) and is few times smaller than that of the former survey.

From the shallow IRAS 1.2 Jy survey conflicting results about the presence of a local void have emerged (*cf* Koranyi & Strauss 1997 and Springel & White 1998), whereas the deeper QDOT-IRAS survey suggests a moderate local void extending out to $\sim 9000 \text{ km s}^{-1}$ (Saunders et al. 1990, Fig. 11). More reliable results may be expected from the larger PSCz survey (e.g., Canavezes et al. 1998) which is under study.

To sum up, some optical redshift surveys (CfA2, CS, ESP) favor a high average normalization, but at the same time hint at a galaxy underdensity in the nearby regions. If we rely on the results coming from these surveys, we should conclude that the local universe is really underdense. But optical and IRAS redshift surveys disagree on the extent, depth, and even real existence of a local underdensity (see §4.4 for further discussions on this point).

4.2 Comparison with previous type-specific galaxy luminosity functions

Our morphological-type dependence of the slope of the optical galaxy LF is in substantial agreement with the results obtained by Marzke et al. (1994b, 1998) from the CfA2 and the SSRS2 samples. The number of Sm-Im objects rises towards the faint end even steeper in our sample than in those two samples (where $\alpha \sim -1.9$). Moreover, the difference between the α -values of E and S0 galaxies (and correspondingly the shortage of low-luminosity E systems) is more marked in our sample than in the CfA2 sample. The difference we find is fully consistent with claims that in the field the transition region between giants and dwarfs is scarcely populated in

the case of ellipticals and better populated in the case of lenticulars, because galaxy bulges tend to weaken with decreasing luminosities and become dominated by disk or lens-like components (see, e.g., the RSA survey of dwarf galaxies by Vader & Chaboyer 1994).

Nevertheless, the dependence of the slope on the morphological type across the whole Hubble morphological sequence is not so striking in our sample as in the Stromlo-APM sample, where the very steep declining of early-type galaxies at the faint end (Loveday et al. 1992) can be ascribed to the incompleteness of early types at high redshift and/or faint magnitudes (as discussed by Marzke et al. 1994b, and Zucca, Pozzetti & Zamorani 1994).

Our analysis establishes that the M_B^* -value of E galaxies is not unusually bright, supporting Marzke et al.'s (1994b) guess that an uncorrected dependence of the Zwicky magnitude scale on surface brightness (e.g., Ichikawa & Fukugita 1992) could have bias the magnitudes of the CfA2 ellipticals.

At middle and faint magnitudes ($M_B > -20$), our spiral LF has an amplitude closer to those of the CfA2 and SSRS2 spiral LFs than to the lower one of the Stromlo-APM late-type LF. At the same magnitudes, the amplitude of our E-S0 LF is close to that of the SSRS2 E-S0 LF.

Interestingly, the spectral type-dependence of the galaxy luminosity function as derived from the LCRS (Bromley et al. 1998 a, b) and the Autofib redshift survey (Heyl et al. 1997) shows a more progressive steepening of the slope from early types to late types than the morphological-type dependence derived in this paper. This is due to the fact that galaxy typing based on the visible morphological appearance is not very tightly correlated with galaxy classification based on spectral schemes (e.g., Kennicutt 1992; Connolly et al. 1995; Zaritsky, Zabludoff & Willick 1995; Katgert et al. 1999) because the latter method is very sensitive to the degree of star formation activity.

Our study lends further support to the fact that blue, gas-rich, late-type spirals and irregulars constitute a steeply rising component of the field galaxy LF. Since these galaxy types dominate in the field at very low luminosities (e.g., Ferguson & Binggeli 1994; Sprayberry et al. 1997), our finding leads us to envisage an upturn of the field LF in the faint end (though not brightwards $M_B \sim -15$). This is consistent with the growing evidence that the field galaxy LF shows an upturn from the Schechter form at faint magnitudes (where magnitude-limited samples become severely censored by surface brightness selection effects). There is straightforward evidence for this faint-end steepening in the CfA2 and ESP data (through an excess of low-luminosity galaxies above the extrapolated Schechter function) and in the counts of the statistical excess of faint APM galaxies seen in projection around Stromlo-APM redshift survey galaxies (Loveday 1997).

4.3 Comparison with previous luminosity functions of group and cluster galaxies

The most recent and extensive analyses of the cluster galaxy LF (Lumsden et al. 1997; Valotto et al. 1997; Gaidos 1997; Rauzy, Adami & Mazure 1998) concur to M_B^* -values brighter by a few tenths of magnitudes than those of the field galaxy LFs mentioned in §4.1 (including our LF and the LFs relying on the same photometric system), whereas there is no agreement on whether the LF slope α of members of (poor and/or rich) clusters significantly differs than that of field galaxy LF (see the conflicting results reached by the above-mentioned authors and also by López-Cruz et al. 1997).

Notably, that tendency is consistent with the environmental dependence of the LF we recognize in §3.3. We find that this dependence is so much weak (at least in the systems of moderate richness which are typical for our sample) that the presence of galaxy systems does not affect significantly the field galaxy LF. This is at variance with some pronounced cluster effects (i.e., a steepening of the slope α in samples containing clusters) reported by Marzke & da Costa (1997) for a small subsample drawn from the SSRS2.

Our results are in line with the claims of some luminosity segregation with density for several galaxy samples (e.g., Park et al. 1994, for the CfA2 galaxies; Giuricin et al. 1995, for the Local Supercluster galaxies; Loveday et al. 1995 for the Stromlo-APM galaxies, Willmer, da Costa & Pellegrini 1998 for the SSRS2 galaxies). This segregation was found to be weaker than morphological segregation with density by Iovino et al. (1993) for the Perseus-Pisces supercluster galaxies and by Domínguez-Tenreiro, Gómez-Flechoso & Martínez (1994) for the CfA1 galaxies, whereas it was considered significant only for early-type luminous galaxies by Hasegawa & Umemura (1993).

Among recent works in which environmental effects on the LF are searched for through a specific comparison between group and field galaxy LFs, an approach similar to ours was followed by Ramella et al. (1999) and Bromley et al. (1988a,b), who divided their respective ESP and LCRS population into field galaxies and galaxy members of groups (or rich groups), identified by means of the *friends of friends* algorithm.

The former authors found that, compared to the LF of field galaxies, the LF of group galaxies exhibits a brighter M^* (by a few tenths of magnitude), which brightens with the increasing richness of groups, whereas the slope α does not change significantly from the field to the groups. Ramella et al.'s (1999) results on changes of M^* are quantitatively compatible with our findings, although the authors claimed a larger (2σ) statistical significance.

On the other hand, the latter authors detected a steepening of the LF slope for galaxies located in regions of higher density. This is caused by a strong environmental effect (in the same

sense) on the LF of early-type galaxies (there is no effect on late types). We must be wary that their steepness-density relation (which we do not find) could reflect some property of the survey (which shows a north-south discrepancy in total and type-dependent LFs) rather than a true environmental effect.

4.4 Relation of the galaxy luminosity function to galaxy counts

The following arguments bearing on the observed galaxy number counts provide a better evidence for an underdensity of the nearby universe than those presented in §4.1. As discussed by several authors (e.g., Driver & Phillipps 1996, and the reviews by Shanks 1989, and Ellis 1997), in the absence of appreciable galaxy evolution, a low normalization of the local galaxy LF, as derived by us, is inconsistent with the high counts which are generally observed already at intermediate optical and near-infrared magnitudes (i.e., roughly at $18 < B < 20$ mag, $13 < K < 15$ mag). The observed counts at these magnitudes would imply a LF normalization greater by a factor ~ 1.5 than that of our LF. A high normalization is customarily assumed in most models of faint galaxy counts, which are generally normalized to the observations in the magnitude intervals quoted above rather than locally. Workers (e.g., Koo, Gronwall & Bruzual 1993; Gronwall & Koo 1995), who claimed that the high bright-end counts can be reasonably fitted by no-evolution or mild evolution models, adopted the view that the local LF is ill-defined.

We deem it more reasonable to explain the underprediction of galaxy counts by conceiving a local underdensity rather than a rapid evolution (Maddox et al. 1990) of the bulk of the galaxy population at the corresponding low redshifts of $z \sim 0.1 - 0.2$ because on theoretical grounds little galaxy evolution is expected up to these redshifts and because very deep redshift surveys which have begun measuring the galaxy LF out to $z \sim 0.5 - 1$ (e.g., Ellis 1997) reveal little evolution in the LF of most of galaxies, except for the late-type low-luminosity population.

Consistently, no-evolution models based on our local LF, but with a normalization roughly 1.5 times our value, would satisfactorily predict the counts of E/S0 and Sabc spirals, but would still underpredict those of late-type spirals and irregulars, for the morphologically classified sample of the faint field galaxies from the HST Medium Deep Survey (Driver, Windhorst & Griffiths 1995; Glazebrook et al. 1995a; Abraham et al. 1996) and the deep HST fields (Odewahn et al. 1996).

As for the compatibility of the observed bright end counts with no-evolution or passive evolution models, recent observations disclaimed the very steep slope of the APM counts reported in the $16 < bj < 19$ mag range by Maddox et al. (1990) — who explained their counts by invoking a large amount of evolution at low redshift — and attributed the steepness of the APM

counts to systematic errors in the non-linear APM photometry. The same effect may account for the low counts of Heydon-Dumbleton, Collins & MacGillivray (1989) and Zucca et al. (1997) at bright COSMOS-*bj* magnitudes. The flatter slopes which fit the bright-end blue counts of recent studies (Weir, Djorgovski & Fayyad 1995; Gardner et al. 1996, Huang et al. 1997, Bertin & Dennefeld 1997) do not largely depart from the expectations of no-evolution models, but always require a higher normalization than our local one.

A more convincing evidence for such a departure comes from the bright-end K-band number counts. At bright magnitudes in the K-band ($10 < K < 17$ mag), where the K-correction is known better than in the optical band and is only a weak function of morphological type, the number counts delineate a slope which is too steep to be consistent with simple no-evolution models (Huang et al. 1997). Detailed models of these counts have explained this effect through a large region of local underdensity of galaxies (by a factor $1.7 - 2.4$), which extends out to $z \sim 0.1 - 0.2$, in the case of no-evolution, or through a combination of some local underdensity and a mild evolution (Phillips & Turner 1998).

The initial steep rise of the radio source counts between fluxes of ~ 10 Jy and ~ 1 Jy (in the 1.4–8.4 GHz frequency range) (e.g., Windhorst et al. 1993) also may suggest the presence of a local underdensity of nearby radiogalaxies, since their strong cosmological evolution is thought not to start until $z \sim 0.3$ (e.g., Condon 1989).

The existence of a large-scale local underdensity implies that local measurements underestimate (overestimate) the true value of Ω_0 (H_0). The local value of H_0 is expected to be higher than the global value (Turner, Cen & Ostriker 1992) by as much as $\sim 20\%$ for $\Omega_0 = 1$ and $\sim 10\%$ for $\Omega_0 = 0.2$, in the case of an underdensity by a factor of 1.5 (see also Shi, Widrow & Dursi 1996). A moderate density contrast (of this order of magnitude or smaller) is lower than the value favored by the above-mentioned models of bright-end K-band galaxy number counts (Phillips & Turner 1998), but it is more consistent with the following observational constraints relative to Hubble flow perturbations. As a matter of fact, from the peculiar velocities of supernovae Zehavi et al. (1998) found a $\sim 7\%$ deviation from the Hubble law consistent with a void of $\sim 20\%$ underdensity surrounded by a dense wall at 7000 km s^{-1} (which roughly coincides with the local Great Walls). This is compatible with the corresponding upper limit ($\sim 7\%$) obtained from the brightest cluster galaxy Hubble diagram (Lauer & Postman 1992; Lauer et al. 1998). Differences between the local and global values of H_0 as large as $\sim 10\%$ on scales of $\sim 30,000 \text{ km s}^{-1}$ are still fully compatible with the limits sets by the measurements of cosmological distances through high redshift supernovae (e.g., Kim et al. 1997), but become only marginally compatible with the limits imposed by the current knowledge of cosmic microwave background anisotropies (e.g.,

Wang, Spergel & Turner 1998).

5 The selection function

The NOG sample, being magnitude-limited, becomes sparser at larger distances due to the increasing loss of galaxies caused by the apparent magnitude cutoff. This effect is quantified by the selection function $S(r)$ which expresses the fraction of galaxies that are expected to satisfy the sample's selection criterion. The incompleteness factor $F(r)$, which expresses the number of galaxies that should have been cataloged for each object present in the sample at a given distance r , is related to $S(r)$ by the expression $F(r) = 1/S(r)$.

The selection function is given by

$$S(r) = f \cdot \begin{cases} \frac{\int_{\max[L_s, L_{\min}(r)]}^{\infty} \Phi_o(L) dL}{\int_{L_s}^{\infty} \Phi_o(L) dL} & \text{if } r > r_s \\ 1 & \text{if } r < r_s \end{cases} \quad (11)$$

where Φ_o is the convolved Schechter LF, $f = 0.8$ is the average completeness level, $L_{\min}(r)$ is the minimum luminosity necessary for a galaxy at distance r (in Mpc) to make it into the sample, and the integral is cut off at the lower limit of $L_s = L_{\min}(r_s)$ corresponding to $M_s = -15.12 + 5 \log h_{75}$ (see §2.2). Through Monte Carlo simulations Santiago et al. (1996)

demonstrated that an unbiased estimate of the galaxy density field can be obtained in the presence of random errors on magnitudes if one uses the convolved LF in eq. 11.

Fig. 7 shows the incompleteness factor F as a function of the distance r (in km s^{-1}) for different velocity field models. The incompleteness factor F is always a smaller in the case of the Mark III multi-attractor model than in the case of the cluster dipole model, but F appears to be little sensitive to the adopted models.

The dependence of the selection function on redshift-space distortions has been addressed analytically by Hamilton (1998) in the case of linear theory. His work is relevant in order to understand the results given in Fig. 7, and then it is mentioned here.

Denoting the redshift-space coordinate by s , the real-space ($F^r(r)$) and redshift-space ($F^s(s)$) incompleteness functions in the LG frame are related as follows (see eq. (4.85) in Hamilton 1998):

$$\ln \left(\frac{F^r(r)}{F^s(s)} \right) = \int_r^{r_{\max}} \frac{d^2 \ln F^s(r')}{d \ln s'^2} \left(\left\langle \frac{v}{r} \right\rangle_{r'} - \left\langle \frac{\hat{\mathbf{r}}}{r} \right\rangle_{r'} \cdot \mathbf{v}^{LG} \right) \frac{dr'}{r'} \quad (12)$$

where $r_{\max}$ is the depth of the sample. Note that averages are not ensemble averages, in which case the real- and redshift-space F would be equal, but are performed over the actual galaxy sample.




Fig. 7.— We plot the incompleteness factor F as a function of galaxy distance r (expressed in km/s) for different peculiar velocity field models.

The second term inside the parenthesis of the right hand side of eq. 12 averages to zero for a full-sky catalog; this highlights a further advantage of the full-sky coverage. The first term in the same parenthesis is the average, at the depth r , of the line-of-sight peculiar velocity v over r . Then, the largest contribution to the differences between the real- and redshift-space

incompletion functions comes from the very nearby universe, both because of the $1/r$ term and because the velocity v is less affected by local fluctuations if the average volume is larger.

With eq 12, considering that the logarithmic derivative is positive for common flux-limited samples, it is possible to understand the behaviors of the different models. As a matter of fact, any model which gives a local overdensity has negative $\langle v/r \rangle_r$ (converging velocities). This is the case for the Mark III multi-attractor model, which has a nearby attractor (the Virgo cluster) and two other attractors (the Great Attractor and the Perseus-Pisces supercluster) at distances of $\sim 4000 - 5000$ km/sec (the distant Shapley concentration gives a significant contribution to the LG motion, but covers a small solid angle). As a consequence, galaxies are brighter (more distant) on average in real space, and the corresponding incompletion factor is consequently smaller.

On the other hand, the cluster dipole model gives a much smoother version of the local velocity field. As a consequence, the local overdensity is low and the incompletion function remains similar to the LG one.

6 Summary and conclusions

We evaluate the field galaxy LF and the related selection function of the nearly complete, magnitude-limited ($B \leq 14$ mag) and volume-limited ($cz < 5500$ km s $^{-1}$) NOG (Nearby Optical Galaxy) sample. Our derivation of the LF is performed using the locations of field and grouped galaxies in real distance space, since we adopt sets of galaxy distances based on different peculiar velocity field models. Our main results can be summarized as follows:

i) The local field galaxy LF is well described by a Schechter function with $\alpha \sim -1.1$, $\Phi^* \sim 0.006$ Mpc $^{-3}$, $M_B^* \sim -20.6$ down to $M_B = -15.1$ ($H_0 = 75$ km s $^{-1}$ Mpc $^{-1}$). The exact values of the Schechter parameters little depend on the adopted velocity field models, since peculiar motion effects are of the order of statistical errors (at most, they cause variations of 0.08 ($\sim 1\sigma$ error) in α and 0.2 mag ($\sim 2\sigma$ error) in M_B^*). Thus, the local field galaxy LF, as well as the intimately related selection function of the NOG sample, are little sensitive to peculiar motions effects. On the other hand, the details of peculiar velocity models have a quite large impact on the local galaxy density on the smallest scales.

ii) Compared to previous local field galaxy LFs, our LF is characterized by a normal (moderately steep) slope α and a M_B^* -value which is, on average, brighter by 0.4 mag than the norm. The last result is explained from the fact that the photometric measures of the NOG galaxies, being total B magnitudes corrected for Galactic absorption, K-dimming, and internal absorption, better represent the true galaxy light.

iii) Our morphological-type dependence of the galaxy LF is in much closer agreement with the results obtained from the CfA2 and SSRS2 samples (Marzke et al. 1994b, 1998) than with those derived from the Stromlo-APM sample (Loveday et al. 1992). The dependence of the galaxy LF from the morphological type appreciably differs from its dependence on the spectral classification as given in the literature.

The LF shape of E-S0 galaxies does not differ significantly from the that of spirals and irregulars. On the other hand, the E galaxies clearly decrease in number towards low luminosities, whereas the number of late-type spirals and irregulars show a very steep rise towards the faint end (with $\alpha \sim -2.3 - -2.4$). This behaviour suggests a similarly fast increase in the number of low-luminosity galaxies in the faint end unexplored in the NOG sample (at $M_B > -15$).

iv) The presence of galaxy systems in the NOG sample does not affect significantly the shape of the the field galaxy LF. As a matter of fact, environmental effects on the total LF are proved to be marginal in the NOG sample. However, the total LF of the galaxy members of the richest systems tends to show a somewhat brighter value of M_B^* (at the 1σ level), which is in line with several earlier claims of galaxy luminosity segregation with density (in particular with Ramella et al.'s (1999) results).

This effect is unrelated to an increased proportion of early-type galaxies in galaxy systems. A luminosity segregation with density could be related to the processes of biased galaxy formation (e.g., Kaiser 1984; Rees 1985) and/or could be induced by late evolutionary processes — such as tidal shocks (e.g., Nogushi & Ishibashi 1986; Sanders et al. 1988), ram pressure effects (Dressler & Gunn 1983), and merging effects (e.g., Barnes & Hernquist 1991) — which can make galaxies more prone to starburst activity whenever they are located in denser environments (e.g., Maia et al. 1994), at least in cases of not extreme environments (Hashimoto et al. 1998, Balogh et al. 1998).

v) Our results supports the fact the normalization of the local galaxy LF is relatively low. Hence, our local galaxy LF underpredicts many observed galaxy number counts (including those of E/S0 galaxies) at relatively bright optical and near-infrared magnitudes (where little galaxy evolution is in general expected and observed). If we do not want to raise serious doubts on the basic aspects of our understanding of galaxy evolution and spectral energy distribution, the most plausible explanation for this underprediction is that the nearby volume we investigate is underdense in galaxies (by a factor of ~ 1.5). However, we are aware of the systematic uncertainties which in general tend to bias low the estimate of the LF normalization factor (e.g., Willmer 1997) and of the appreciable sensitivity of this factor to density anomalies within local volumes.

Notably, uncertainties related to the choice of the global LF normalization have a much larger

impact on the census of light in the universe (hence, on the determination of the critical mass-to-light ratio) than uncertainties in the light contribution of uncataloged LSB galaxies (see Dalcanton, Spergel & Summers 1997b for a dissenting view).

In conclusion, large observational efforts (e.g., redshift surveys) are still needed to reliably establish the existence, density contrast, and size of a possible local void, which, as suggested by some authors, can be larger than the $\sim 5000 \text{ km s}^{-1}$ radius region examined in this paper.

Acknowledgments

We wish to thank S. Bardelli, M. Girardi, G. Giudice, S. J. Maddox, F. Mardirossian, M. Mezzetti, M. Ramella for interesting conversations and also M. A. Strauss for helpful comments.

In this work we have made use of the Lyon-Meudon Extragalactic Database (LEDa) supplied by the LEDa team at the CRAL-Observatoire de Lyon (France).

P. M. has been supported by the EC contract ERB FMB ICT961709. B. C. has been supported by a CIRAC fellowship. C. M. thanks SISSA for its kind hospitality.

This work has been partially supported by the Italian Ministry of University, Scientific and Technological Research (MURST) and by the Italian Space Agency (ASI).

REFERENCES

- Abraham, R. G. et al. 1996 *ApJS*, 107, 1.
Bahcall, N. A., Lubin, L. M. & Dorman, V. 1995, *ApJ*, 447, L81.
Baker, J. E., Davis, M., Strauss, M. A., Lahav, O. & Santiago, B. X. 1998, *ApJ*, 508, 6.
Balogh M. L. et al. 1998, *ApJ*, 504, L75.
Barnes, J. E. & Hernquist, L. E. 1991, *ApJ*, 370, L65.
Bertin, E. & Dennefeld, M. 1997, *A&A*, 317, 43.
Binggeli, B., Sandage, A. & Tammann, G. A. 1988, *ARA&A*, 26, 509.
Biviano, A. et al. 1995, *A&A*, 297, 610.
Branchini, E. & Plionis, M. 1996, *ApJ*, 460, 569.
Bristow, P. D. & Phillipps, S. 1994, *MNRAS*, 267, 13.
Bromley, B. C., Press, W. H., Lin, H. & Kirschner, R. P. 1998a, *ApJ*, 505, 25.
Bromley, B. C., Press, W. H., Lin, H. & Kirschner, R. P. 1998b, preprint astro-ph 9805197.
Burstein, D. & Heiles, C. 1978a, *ApJ*, 19, 69.
Burstein, D. & Heiles, C. 1978b, *ApJ*, 225, 40.
Burstein, D. & Heiles, C. 1982, *AJ*, 87, 1165.
Burstein, D. & Heiles, C. 1984, *ApJS*, 54, 33.
Buta, R. & Williams, K. L. 1995, *AJ*, 109, 543.
Canavezes, A. et al. 1998, *MNRAS*, 297, 777.
Colless, M. 1989, *MNRAS*, 237, 799.
Colless, M. 1997, in *Wide-Field Spectroscopy*, ed. E. Kontizas et al. (Dordrecht: Kluwer Academic Press), 227.
Condon, J. J. 1989, *ApJ*, 338, 13.
Connolly, A. J., Szalay, A. S., Bershad, M. A., Kinney, A. L. & Calzetti, D. 1995, *AJ*, 110, 1071.
Cowie, L. L., Songaila, A., Hu, E. M. & Cohen, J. G. 1996, *AJ*, 112, 839.
da Costa, L. N. et al. 1994, *ApJ*, 424, L1.
Dalcanton, J. J. 1998, *ApJ*, 495, 251.
Dalcanton, J. J., Spergel, D. N., Gunn, J. E., Schmidt, M. & Schneider, D. P. 1997a, *AJ*, 114, 635.
Dalcanton, J. J., Spergel, D. N. & Summers, F. J. 1997b, *ApJ*, 482, 659.
David, L. P., Jones, C. & Forman, W. 1995, *ApJ*, 445, 578.
Davies, J., Disney, M. J. & Phillips, S. 1989, *MNRAS*, 239, 939.
Davis, M. & Huchra, J. P. 1982, *ApJ*, 254, 437.
Dekel, A., Burstein, D. & White, S. 1997, in *Critical Dialogues in Cosmology*, ed. N. Turok (Singapore: World Scientific).
de Lapparent, V., Geller, M. J. & Huchra, J. P. 1989, *ApJ*, 343, 1.
de Vaucouleurs, G., de Vaucouleurs, A., Corwin, H. G. 1976, *Second Reference Catalogue of Bright Galaxies* (Austin: Univ. of Texas Press).
de Vaucouleurs, G. et al. 1991, *Third Reference Catalogue of Bright Galaxies* (New York: Springer Verlag).
Disney, M. J. & Phillips, S. 1985, *MNRAS*, 216,

- 53,
Dominguez-Tenreiro, R., Gómez-Flechoso, M. A. & Martínez, V. J. 1994, *ApJ*, 424, 42.
Dressler, A. & Gunn, J. E. 1983, *ApJ*, 270, 7.
Driver, S. P. & Phillipps, S. 1996, *ApJ*, 469, 529.
Driver, S. P., Windhorst, R. A. & Griffiths, R. E. 1995, *ApJ*, 453, 48.
Efstathiou, G., Ellis, R. S. & Peterson, B. S. 1988, *MNRAS*, 232, 431.
Ellis, R. S. 1997, *ARA&A*, 35, 389.
Ellis, R. S., Colless, M., Broadhurst, T., Heyl, J. & Glazebrook, K. 1996, *MNRAS*, 280, 235.
Ferguson, H. C. & Binggeli, B. 1994, *ARA&A*, 6, 67.
Gaidos, E. J. 1997, *AJ*, 113, 117.
Garcia, A. M. 1993, *A&AS*, 100, 47.
Garcia, A. M., Paturel, G., Bottinelli, L. & Gougouenheim, L. 1993, *A&AS*, 98, 7.
Gardner, J. P., Sharples, R. M., Carrasco, B. E. & Frenk, C. S. 1996, *MNRAS*, 282, L1.
Gardner, J. P., Sharples, R. M., Frenk, C. S. & Carrasco, B. E. 1997, *ApJ*, 480, L99.
Geller, M. J., Kurtz, M., Wegner, G., Thorstensen, J. R., Fabricant, D. G. & Marzke, R. O. 1997, *AJ*, 114, 2205.
Girardi, M. et al. 1999, in preparation.
Giuricin, G., F. Limboz Tektunali, P. Monaco, F. Mardirossian & M. Mezzetti 1995, *ApJ*, 450, 41.
Glazebrook, K., Ellis, R. S., Santiago, B. & Griffiths, R. E. 1995a, *MNRAS*, 275, L19.
Glazebrook, K., Peacock, J. A., L. Miller & Collins, C. A. 1995b, *MNRAS*, 275, 169.
Gronwall, C. & Koo, D. C. 1995, *ApJ*, 440, L1.
Gudehus, D. 1995, *A&A*, 302, 21.
Hamilton, A. J. S. 1998, in *Ringberg Workshop on Large-Scale Structure*, ed. D. Hamilton (Dordrecht: Kluwer Academic Press), 185.
Hasegawa, T. & Umemura, M. 1993, *MNRAS*, 263, 191.
Hashimoto, Y., Oemler, A., Jr., Lin, H. & Tucker, D. L. 1998, *ApJ*, 499, 589.
Heydon-Dumbleton, N. H., Collins, C. A. & MacGillivray, H. T. 1989, *MNRAS*, 238, 379.
Heyl, J., Colless, M., Ellis, R. S. & Broadhurst, T. 1997, *MNRAS*, 285, 613.
Huang, J.-S., Cowie, L. L., Gardner, J. P., Hu, E. M., Songaila, A. & Wainscoat, R. J. 1997, *ApJ*, 476, 12.
Huchra, J. P. & Geller, M. 1982, *ApJ*, 257, 423.
Hudson, M. J. 1993, *MNRAS*, 265, 43.
Ichikawa, T. & Fukugita, M. 1992, *ApJ*, 394, 61.
Impey, C. D. & Bothun, G. D. 1997, *ARA&A*, 35, 267.
Impey, C. D., Bothun, G. D. & Malin, D. F., 1988, *ApJ*, 330, 534.
Iovino, A., Giovanelli, R., Haynes, M., Chincarini, G., Guzzo, L. 1993, *MNRAS*, 265, 21.
Irwin, M. J., Davies, J. I., Disney, M. J. & Phillipps, S. 1990, *MNRAS*, 245, 289.
Kaiser, N. 1984, *ApJ*, 284, L9.
Katgert, P. et al. 1999, in *Observational Cosmology: The Development of Galaxy Systems*, ed. G. Giuricin, M. Mezzetti & P. Salucci (San Francisco: ASP Conference Series), in preparation.
Kennicutt, R. C., Jr. 1992, *ApJ*, 388, 310.
Kim, A. G. 1997, *ApJ*, 476, L63.
Koo, D. C., Gronwall, C. & Bruzual, A. G. 1993, *ApJ*, 415, L21.
Koranyi, D. M. & Strauss, M. A. 1997, *ApJ*, 477, 36.
Lauer, T. R. & Postman, M. 1992, *ApJ*, 400, L47.
Lauer, T. R., Tonry, J. L., Postman, M., Ajhar, E. A. & Hotlzman, J. A. 1998, *ApJ*, 499, 577.
Lin, H. et al. 1996, *ApJ*, 464, 60.
López-Cruz, O., Yee, H. K. C., Brown, J. P., Jones, C. & Forman, W. 1997, *ApJ*, 475, L97.
Loveday, J. 1997, *ApJ*, 489, 29.
Loveday, J., Maddox, S. J., Efstathiou, G. & Peterson, B. A., 1995, *ApJ*, 442, 457.
Loveday, J., Peterson, B. A., Efstathiou, G. & Maddox, S. J. 1992, *ApJ*, 390, 338.
Lumsden, S. L., Collins, C. A., Nichol, R. C., Eke, V. R. & Guzzo, L. 1997, *MNRAS*, 290, 119.
Maddox, S. J. 1998, in *The Young Universe: Galaxy Formation and Evolution at Intermediate and High Redshift*, ed. S. D'Odorico, A. Fontana & E. Giallongo, E. (San Francisco: ASP Conference Series), 198.
Maddox, S. J., Sutherland, W. J., Efstathiou, G., Loveday, J. & Peterson, B. A. 1990, *MNRAS*, 247, 1P.
Maia, M. A. G., Pastoriza, M. G., Bica, E. & Dottori, H. 1994, *ApJS*, 93, 425.
Marinoni, C., Monaco, P., Giuricin, G. & Costantini, C. 1998a, in *The Young Universe: Galaxy Formation and Evolution at Intermediate and High Redshift*, ed. S. D'Odorico, A. Fontana & E. Giallongo (San Francisco: ASP Conference Series), 202.
Marinoni, C., Monaco, P., Giuricin, G. & Costantini, C. 1998b, *ApJ*, 505, 484 (Paper I).
Marzke, R. O. & da Costa, L. N. 1997, *AJ*, 113, 185.
Marzke, R. O., da Costa, L. N., Pellegrini, P. S., Willmer, C. N. A. & Geller, M. 1998, *ApJ*, 503, 617.
Marzke, R. O., Geller, M., Huchra, J. P. & Corwin, Jr., H. G. 1994b, *AJ*, 108, 437.
Marzke, R. O., Huchra, J. P. & Geller, M. 1994a, *ApJ*, 428, 43.
Metcalf, N., Fong, R. & Shanks, T. 1995, *MNRAS*, 274, 769.
Mobasher, B., Sharples, R. M. & Ellis, R. S. 1993, *MNRAS*, 263, 560.
Noguchi, M. & Ishibashi, S. 1986, *MNRAS*, 219, 305.
Odewahn, S. C., Windhorst, R. A., Driver, S. P. & Keel, W. C. 1996, *ApJ*, 472, L13.
Park, C., Vogeley, M. S., Geller, M. J. & Huchra, J. P. 1994, *ApJ*, 431, 569.
Paturel, G., Bottinelli, L. & Gougouenheim, L. 1994, *A&A*, 286, 768.
Pence, W. D. 1976, *ApJ*, 203, 39.
Phillipps, S., Davies, J. L. & Disney, M. J. 1990, *MNRAS*, 242, 235.
Phillips, L. A. & Turner, E. L. 1998, preprint, astro-ph/9802352.
Ramella, M. et al. 1999, *A&A*, in press.
Ratcliffe, A., Shanks, T., Parker, Q. A. & Fong, R. 1998, *MNRAS*, 293, 197.

- Rauzy, S., Adami, C. & Mazure, A. 1998, *A&A*, 337, 31.
- Rees, M. J. 1985, *MNRAS*, 213, 75P.
- Rousseau, J., Di Nella, H., Paturel, G. & Petit, C. 1996, *MNRAS*, 282, 144.
- Sandage, A. & Tammann, G. A. 1981, *A Revised Shapley-Ames Catalog of Bright Galaxies* (Washington: Carnegie Inst. of Washington) (RSA).
- Sanders, D. B. et al. 1988, *ApJ*, 325, 74.
- Santiago, B. X., Strauss, M. A., Lahav, O. et al. 1995, *ApJ*, 446, 457.
- Santiago, B. X., Strauss, M. A., Lahav, O. et al. 1996, *ApJ*, 461, 38.
- Saunders, W. et al. 1990, *MNRAS*, 242, 318.
- Schechter, P. L. 1976, *ApJ*, 203, 297.
- Shanks, T. 1989, in *The Extragalactic Background Light*, ed. S. C. Bowyer & C. Leinert (Dordrecht: Kluwer Academic Publ.), 269.
- Shanks, T., Stevenson, P. R. F., Fong, R., MacGillivray, H. T. 1984, *MNRAS*, 206, 767.
- Shi, X., Widrow, L. M. & Dursi, L. J. 1996, *MNRAS*, 281, 565.
- Sprayberry, D., Impey, C. D., Irwin, M. J. & Bothun, G. D. 1997, *ApJ*, 481, 104.
- Springel, V. & White, S. D. M. 1998, *MNRAS*, 298, 143.
- Szokoly, G. P., Subbarao, M. U., Connolly, A. J. & Mobasher, B. 1998, *ApJ*, 492, 452.
- Takamiya, M., Kron, R. G. & Kron, G. E. 1995, *AJ*, 110, 1083.
- Tully, R. B. 1987, *ApJ*, 321, 280.
- Tully, R. B. & Fouqué, P. 1985, *ApJS*, 58, 67.
- Turner, E. L. 1979, *ApJ*, 231, 645.
- Turner, E. L., Cen, R. & Ostriker, J. P. 1992, *AJ*, 103, 1427.
- Vader, J. P. & Chaboyer, B. 1994, *AJ*, 108, 1209.
- Valotto, C. A., Nicotra, M. A., Muriel, H. & Lambas, D. G. 1997, *ApJ*, 479, 90.
- Wang, Y., Spergel, D. N. & Turner, E. L., 1998, *ApJ*, 498, 1.
- Weir, N., Djorgovski, S. & Fayyad, U. M. 1995, *AJ*, 110, 1.
- Willick, J. A., Courteau, S., Faber, S. M., Burstein, D., Dekel, A., Strauss, M. A. 1997, *ApJS*, 109, 333.
- Willmer, C. N. A. 1997, *AJ*, 114, 898.
- Willmer, C. N. A., da Costa L. N. & Pellegrini, P. S. 1998, *AJ*, 115, 869.
- Wilson, G., Smail, I., Frenk, C. S., Ellis, R. S. & Couch, W. J. 1997, *MNRAS*, 284, 915.
- Windhorst, R. A., Fomalont, E. B., Partridge, R. B. & Lowenthal, J. D. 1993, *ApJ*, 405, 498.
- Yahil, A., Strauss, M. A., Davis, M., & Huchra J, P. 1991, *ApJ*, 372, 380.
- Zaritsky, D., Zabludoff, A. I. & Willick, J. A. 1995, *AJ*, 110, 1602.
- Zehavi, I., Riess, A. G., Kirschner, R. P. & Dekel, A., 1998, *ApJ*, 503, 483.
- Zucca, E., Pozzetti, L. & Zamorani, G. 1994, *MNRAS*, 269, 953.
- Zucca, E. et al., 1997, *A&A*, 326, 477.